

A Strong threshold for the size of random caps to cover a sphere.

by

Bhupendra Gupta ¹

Faculty of Engineering and Sciences,
Indian Institute of Information Technology (DM)-Jabalpur, India.

Abstract

In this article, we consider ' N 'spherical caps of area $4\pi p$ were uniformly distributed over the surface of a unit sphere. We are giving the strong threshold function for the size of random caps to cover the surface of a unit sphere. We have shown that for large N , if $\frac{Np}{\log N} > 1/2$ the surface of sphere is completely covered by the N caps almost surely, and if $\frac{Np}{\log N} \leq 1/2$ a partition of the surface of sphere is remains uncovered by the N caps almost surely.

AMS 2000 subject classifications: 05C80, 91D30.

Keywords: Coverage Problem, Random Caps, Threshold Function.

¹Corresponding Author. email:gupta.bhupendra@gmail.com, bhupen@iiitdm.in

1 Introduction.

Let $V_1, V_2, \dots, V_N$ be the spherical caps on the surface of a unit sphere with their centers $v_1, v_2, \dots, v_N$ respectively, on the surface of a unit sphere. Also let $v_1, v_2, \dots, v_N$ are independently and uniformly distributed on the surface of a unit sphere. H. Maehara, [2] gives the threshold function $p_0(N) = \frac{\log N}{N}$ for the coverage of the surface of a unit sphere. H. Maehara, proves that for $\frac{p(N) \cdot N}{\log N} < 1$, probability that N spherical caps cover the entire surface of the unit sphere is converges to 0, and for $\frac{p(N) \cdot N}{\log N} > 1$, probability that each point of the sphere is covered by n caps is converges to 1. Since both of these are convergence in probability sense, the threshold $p_0(N)$ is a weak threshold. Also in article [2], instead of exact bounds author use loose approximations. Due to these approximations threshold suggested in [2] is different from threshold suggested in this article.

Now using the same model and notations as in H. Maehara, [2], we are giving the strong threshold function for the coverage of the surface of a unit sphere.

2 Basic Model and Definitions.

Here, we recall the same model as it is given by H. Maehara, [2]. We made some modification in the language for making the things more clear.

Let ‘ S ’ be the surface area of a unit sphere in 3—dimensional space. Let $V_1, V_2, \dots$, be the spherical caps on the surface of a unit sphere with their centers $v_1, v_2, \dots$, respectively, and uniformly distributed on the surface of a unit sphere. The area of a spherical cap of angular distance (angular radius) ‘ a ’ is $2\pi(1 - \cos(a)) = 4\pi \sin^2(a/2)$.

Let ‘ p ’ be the probability that any point on the surface of unit sphere covered by a

specified spherical cap of angular distance (angular radius) ‘ a ’. Then

$$p := \frac{\text{Area of a spherical cap of angular distance 'a'}}{\text{Surface area of unit sphere}} = \frac{4\pi \sin^2(a/2)}{4\pi} = \sin^2(a/2) \quad (2.1)$$

Let there are ‘ N ’ random caps of angular distance ‘ a ’ on the surface ‘ S ’ of unit sphere. Let $U_0(N, p)$, be the set of those points which remains uncovered by ‘ N ’ spherical caps and $u_0(N, p)$ be the proportion of the area covered by $U_0(N, p)$, i.e., $u_0(N, p)$ be the proportion of the area which remains uncovered by ‘ N ’ spherical caps:

$$u_0(N, p) := \frac{\{\text{the area of } U_0(N, p)\}}{4\pi}. \quad (2.2)$$

Then

$$E(u_0(N, p)) = \frac{1}{4\pi} \int_S P[x \in U_0(N, p)] dx. \quad (2.3)$$

Now consider,

$$\begin{aligned} P[x \in U_0(N, p)] &= P[x \text{ is remains uncovered}] \\ &= \prod_{i=1}^N (1 - P[x \in V_i,]) \\ &= (1 - p)^N. \end{aligned} \quad (2.4)$$

Hence, from (2.3), we have

$$E(u_0(N, p)) = (1 - p)^N. \quad (2.5)$$

Similarly, as in H. Maehara, [2], we have

$$E(u_0^2(N, p)) = \frac{1}{16\pi^2} \int_S \int_S P[x, y \in U_0(N, p)] dx = \frac{1}{4\pi} \int_S P[x_0, y \in U_0(N, p)] dy, \quad (2.6)$$

where x_0 is a fixed point on S . Let x_0 and y subtend an angle ‘ θ ’ at the center of sphere.

Then

$$P[x_0, y \in U_0(N, p)] = (1 - (2p - q(\theta)))^N, \quad (2.7)$$

where, $q(\theta)$ be the area of intersection between two spherical caps of angular distance ‘ a ’. Substituting the above probability in (2.6), we get

$$E(u_0^2(N, p)) = \frac{1}{4\pi} \int_S (1 - (2p - q(\theta)))^N dy.$$

Since points x_0 and y subtend an angle between θ and $\theta + d\theta$ at the center of the sphere.

Then

$$E(u_0^2(N, p)) = \int_0^\pi (1 - (2p - q(\theta)))^N (1/2) \sin(\theta) d\theta. \quad (2.8)$$

Since $q(\theta) = 0$ for $\theta > 2a$,

$$\begin{aligned} E(u_0^2(N, p)) &< \int_0^{2a} (1 - p)^N (1/2) \sin(\theta) d\theta + \int_{2a}^\pi (1 - 2p)^N (1/2) \sin(\theta) d\theta \\ &< (1 - p)^N [-(1/2) \cos(\theta)]_0^{2a} + (1 - 2p)^N \\ &= (1 - p)^N \frac{1 - \cos(2a)}{2} + (1 - 2p)^N. \end{aligned} \quad (2.9)$$

Using (2.5), in (2.9), we have

$$\begin{aligned} E(u_0^2(N, p)) &< (1 - p)^N \frac{1 - \cos(2a)}{2} + (1 - 2p)^N \\ &= 4p(1 - p)^{N+1} + (1 - 2p)^N, \end{aligned} \quad (2.10)$$

since $\frac{1 - \cos(2a)}{2} = 4p(1 - p)$.

Now for the lower bound of $E(u_0^2(N, p))$, from (2.8) we have

$$E(u_0^2(N, p)) > \int_{2a}^\pi (1 - (2p - q(\theta)))^N (1/2) \sin(\theta) d\theta, \quad (2.11)$$

since $q(\theta) = 0$ for $\theta > 2a$.

$$\begin{aligned} E(u_0^2(N, p)) &> \frac{(1 - 2p)^N}{2} \int_{2a}^\pi \sin(\theta) d\theta \\ &= \frac{(1 - 2p)^N}{2} [-\cos(\theta)]_{2a}^\pi \\ &= (1 - 2p)^N \frac{\cos(2a) + 1}{2} \\ &= (1 - 2p)^N (1 - 4p(1 - p)), \end{aligned} \quad (2.12)$$

since $\frac{\cos(2a) + 1}{2} = 1 - 4p(1 - p)$.

Let Θ be a fixed monotone property.

Definition 1 A function $\delta_\Theta(c) : Z^+ \rightarrow R^+$ is a strong threshold function for Θ if the following is true for every fixed $\epsilon > 0$,

- if $P[\delta_\Theta(c - \epsilon) \in \Theta] = 1 - o(1)$, and
- if $P[\delta_\Theta(c + \epsilon) \in \Theta] = o(1)$,

where ‘ c ’ is some constant. □

3 Main Result.

Theorem 3.1 Let $p = \frac{c \log N}{N}$, then for $c > \frac{1}{2}$, we have the surface of unit sphere is completely covered by ‘ N ’ spherical caps, i.e.,

$$U_0(N, p) = \phi, \quad \text{almost surely,}$$

and for $c \leq \frac{1}{2}$, surface of unit sphere is not completely covered by ‘ N ’ spherical caps, i.e.,

$$U_0(N, p) \neq \phi, \quad \text{almost surely.}$$

Proof. For arbitrary small ϵ . We have,

$$P[U_0(N, p) \neq \phi] \simeq P[|u_0(N, p)| \geq \epsilon]. \quad (3.13)$$

From the Markov’s inequality, we have

$$\begin{aligned} P[|u_0(N, p)| \geq \epsilon] &\leq \frac{E[u_0^2(N, p)]}{\epsilon^2} \\ &< \frac{1}{\epsilon^2} \left(4p(1-p)^{N+1} + (1-2p)^N \right), \end{aligned} \quad (3.14)$$

using the upper bound of $E[u_0^2(N, p)]$ from (2.10). Now taking $p = \frac{c \log N}{N}$, where c is some constant. Then

$$P[|u_0(N, p)| \geq \epsilon] < \frac{1}{\epsilon^2} \left(\frac{4c \log N}{N} \left(1 - \frac{c \log N}{N} \right)^{N+1} + \left(1 - \frac{2c \log N}{N} \right)^N \right)$$

$$\begin{aligned}
&< \frac{1}{\epsilon^2} \left(\frac{4c \log N}{N} \left(1 - \frac{c \log N}{N} \right) e^{-c \log N} + e^{-2c \log N} \right) \\
&< \frac{1}{\epsilon^2} \left(\frac{4c \log N}{N^{1+c}} + \frac{1}{N^{2c}} \right).
\end{aligned} \tag{3.15}$$

If $c > 1/2$, the above probability is summable, i.e.,

$$\sum_{N=0}^{\infty} P[|u_0(N, p)| \geq \epsilon] < \infty,$$

and hence from (3.13), we have

$$\sum_{N=0}^{\infty} P[U_0(N, p) \neq \phi] < \infty.$$

Then by the Borel-Cantelli's Lemma, we have

$$P[U_0(N, p) \neq \phi, \quad i.o.] = 0.$$

Thus, the set $U_0(N, p) \neq \phi$ for only finitely many times, i.e., eventually $U_0(N, p) = \phi$ happens infinitely times with probability 1. Hence for $c > \frac{1}{2}$, we have

$$U_0(N, p) = \phi, \quad \text{almost surely.}$$

Now, by the lower bound of Chebyshev's inequality (Page 55, Shirayev [3].), we have

$$P[u_0(N, p) \geq \epsilon] \geq \frac{E[u_0(N, p)^2] - \epsilon^2}{16\pi^2}, \tag{3.16}$$

since $u_0(N, p) \geq 0$ and $|u_0(N, p)| \leq 4\pi$. Now using the lower bound of $E[u_0(N, p)^2]$ from (2.12) and taking $\epsilon = o\left(\frac{1}{N}\right)$, we have

$$\begin{aligned}
P[u_0(N, p) \geq \epsilon] &\geq \frac{(1 - 2p)^N (1 - 4p(1 - p)) - \epsilon^2}{16\pi^2} \\
&\geq C_1(1 - 2p)^N (1 - 4p(1 - p)) - C_2,
\end{aligned} \tag{3.17}$$

where $C_1 = \frac{1}{16\pi^2}$ and $C_2 = \frac{\epsilon^2}{16\pi^2}$.

Substituting $p = \frac{c \log N}{N}$, in the above expression we get

$$\begin{aligned} P[u_0(N, p) \geq \epsilon] &\geq C_1 \left(1 - \frac{2c \log N}{N}\right)^N \left(1 - \frac{4c \log N}{N} \left(1 - \frac{c \log N}{N}\right)\right) - C_2 \\ &\geq C_3 e^{-2c \log N} = \frac{C_3}{N^{2c}}, \end{aligned} \quad (3.18)$$

where C_3 is some constant. If we take $c \leq 1/2$, then the probability (3.18), is not summable with respect to N , i.e.,

$$\sum_{N=1}^{\infty} P[u_0(N, p) \geq \epsilon] = \infty.$$

Then by the Borel-Cantelli's Lemma, we have

$$P[u_0(N, p) \geq \epsilon, \quad i.o.] = 1,$$

since $u_0(N, p)$ are independent. Thus $u_0(N, p) \geq \epsilon$ happens infinitely many time with probability 1. Hence for $c \leq \frac{1}{2}$, we have

$$u_0(N, p) \geq \epsilon, \quad \text{almost surely.}$$

This implies for $c \leq 1/2$, we have $U_0(N, p) \neq \phi$ almost surely. $\square$

References

- [1] Y.S.Chow, H.Teicher(2004), Probablity Theory, thired edition, *Springer Text in Statistics*.
- [2] H. Maehara, (1988), A threshold for the size of random caps to cover a sphere, *Annal of the Institute of Statistical Mathematics*, Vol. 40, No.4, 665 - 670.
- [3] Shirayayev, A.N.(1984), Probability, second edition, *Springer-Verlag, New York Inc*.
- [4] M. Karonski, E.R.Scheinerman and K.B.Singer-Cohen, (1999), On random inter-section graphs: the subgraph problem, *Combinatorics, Probability and Computing*, Vol. 8, 131-159.